

[CASE REPORT AND LITERATURE REVIEW]

Amyopathic Dermatomyositis Complicated by Pneumomediastinum

^aRANDY TANG, RN, BSN, BSE, MD; ^bCHRISTIAN R. MILLETT, MD; ^cJUSTIN J. GREEN, MD

^aUMDNJ—Robert Wood Johnson Medical School, Camden, New Jersey; ^bDermatology Center, Bethesda, Maryland;

^cCooper University Hospital, Camden, New Jersey

ABSTRACT

Dermatomyositis is an inflammatory disease of unclear etiology with characteristic cutaneous and musculoskeletal findings. Amyopathic dermatomyositis is a subtype without musculoskeletal involvement. Many cases of dermatomyositis are associated with underlying malignancy, but pulmonary manifestations can also be seen, the most common of which is interstitial lung disease. Pneumomediastinum is a rare complication that is important for clinicians to recognize, as it may be fatal if left untreated. The sudden onset of facial edema and shortness of breath in the setting of dermatomyositis should raise the suspicion of this condition. (*J Clin Aesthet Dermatol.* 2013;6(3):40–43.)

A 52-year-old Caucasian woman presented to the authors' clinic with a six-month history of fatigue, hair loss, and joint stiffness of the ankles, wrists, and knees. On physical exam, the patient was noted to have periorbital erythema with profound edema (Figure 1). Confluent and violaceous erythema was noted over the upper chest, and violaceous, ulcerated plaques were present on the metacarpalphalangeal joints, distal interphalangeal joints, and elbows bilaterally (Figure 2). A punch biopsy of the left elbow plaque was performed at the time.

The patient was started on prednisone 60mg once daily for a presumptive diagnosis of dermatomyositis (DM), and a comprehensive laboratory evaluation was ordered. The biopsy results were consistent with DM, and anti-Jo-1 antibody levels were normal. As her creatine kinase and aldolase levels were also normal, a diagnosis of amyopathic dermatomyositis (ADM) was made. At the next office visit, the patient showed no signs of clinical improvement, so hydroxychloroquine 200mg twice daily and clobetasol ointment were added. This combination led to significant improvement in the patient's cutaneous lesions.

One year after initial presentation, the patient came to the authors' clinic with an acute episode of shortness of breath as well as facial and neck edema. Crepitus was appreciated on physical exam. She was sent to the nearest emergency room where a computed tomography scan revealed bilateral pneumomediastinum (PnM) with

subcutaneous emphysema. Bilateral chest tubes were placed at the time and cyclosporine therapy was initiated. The tubes were removed and the patient was discharged one week later after significant improvement in her condition.

DISCUSSION

Dermatomyositis is an inflammatory disease of unclear etiology characterized by proximal muscle weakness and cutaneous manifestations, including erythema of the eyelids with periorbital edema (heliotrope rash), erythematous to violaceous papules over the extensor aspects of the interphalangeal joints (Gottron's papules) as well as elbows and knees (Gottron's sign), hyperpigmentation with hyperkeratosis and fissuring of the hands (mechanic's hands), and erythema of the neck and upper chest in a v-shaped distribution (shawl sign). Amyopathic dermatomyositis is a subtype of DM in which patients present with typical cutaneous findings without associated muscle weakness or elevated muscle enzyme levels. The etiology of DM is a vasculopathic process with deposition of immune complexes on the endothelium of the capillaries in the skin as well as in the muscle.^{1,2} Severe cases of DM may therefore present with ulcerative or even necrotic skin lesions.

It is well established that DM is associated with malignancy.^{3,4} In some instances, the cutaneous

DISCLOSURE: The authors report no relevant conflicts of interest.

ADDRESS CORRESPONDENCE TO: Christian Millett, MD, Dermatology Center, 6410 Rockledge Drive, Suite 201, Bethesda, MD 20817;
E-mail: millett.christian@gmail.com



Figure 1. Significant facial and neck swelling of acute onset



Figure 2. Erythematous, ulcerated papules overlying the MCP, PIP, and DIP joints of the hands

manifestations of DM are the first indication of an underlying malignancy. Additionally, it is estimated that approximately 50 percent of patients with DM will develop some type of pulmonary manifestation of their disease.⁵ These include interstitial lung disease (ILD), aspiration pneumonia from esophageal dysfunction, pulmonary edema secondary to DM-induced cardiomyopathy, infectious pneumonia from drug-induced immunosuppression, pulmonary fibrosis secondary to methotrexate or cyclophosphamide therapy, or hypoventilation secondary to inflammation of the respiratory muscles. The most common pulmonary complication of DM is interstitial lung disease, and nonspecific interstitial pneumonitis (NSIP) accounts for 80 percent of these cases.⁶ Patients with DM-associated ILD are more likely to have the anti-aminoacyl transfer RNA synthetase antibody (Anti-Jo-1), which has been termed the anti-synthetase syndrome. This antibody is present in 70 percent of DM patients with ILD, but only 20 percent of those without ILD.⁷⁻⁹ However, the antibody is typically absent in patients with ADM-associated ILD.¹⁰ Morganroth et al¹¹ reported that there was no difference in ILD prevalence between DM and ADM. Mukae et al¹² compared patients with DM-associated ILD and those with ADM-associated ILD and found a 45-percent mortality rate among those with DM as compared to a six-percent mortality rate for those with ADM.

Pneumomediastinum is a rare complication of DM that can be fatal if left unrecognized. Bradley¹³ described the first case of PnM in an adult patient with DM, and since then, there have been several dozen reported cases, some in association with DM, others in association with ADM. Several mechanisms have been proposed to account for the origin of PnM in DM patients. One hypothesis is that PnM occurs when raised intra-alveolar pressure ruptures previously damaged alveoli in the lung or ruptures a

subpleural cyst that developed from interstitial fibrosis.¹⁴ A second hypothesis presumes that underlying pulmonary vasculitis leads to rupture of airway lesions,^{15,16} while a third suggests that corticosteroids in the course of DM treatment weakens alveolar walls, predisposing to rupture.¹⁷⁻¹⁹ Each of the above hypothesized mechanisms may contribute or act synergistically to increase the risk of PnM formation. The risk factors that predispose patients with DM to develop PnM therefore include the presence of interstitial lung disease (ILD), cutaneous vasculopathy, and a history of previous systemic corticosteroid treatment. Interestingly, younger age and absent creatine kinase elevation are risk factors as well.²⁰

The treatments of choice for PnM are immunosuppressive agents, such as cyclosporine,^{16,21,22,27} methotrexate,^{17,18} and cyclophosphamide.²³ These should also be first-line treatments in patients with underlying ADM-associated ILD.²⁴ Other therapeutic options include the use of corticosteroids²⁵ and rituximab.²⁶ A review of previous cases highlights the variety of therapies that have been utilized (Table 1).

Although PnM may be fatal, many cases will self-resolve. Those cases reported in the literature that had a fatal outcome have typically occurred in individuals with concurrent ILD. The ultimate cause of death in these cases may not be from the PnM itself, but rather from respiratory failure secondary to progressive ILD. Hence, Yoshida et al¹⁷ suggest that PnM in DM is not a prognostic factor. Rather, the presence and severity of ILD dictates the prognosis.^{17,27}

Although uncommon, many physicians encounter patients with dermatomyositis. The sudden onset of facial edema and shortness of breath in the setting of DM should raise the suspicion of pneumomediastinum, as prompt recognition may be lifesaving. Other diagnoses in the differential include an allergic reaction, anaphylaxis, angioedema, and superior vena cava (SVC) syndrome.

TABLE 1. Previous cases and the variety of therapies that have been utilized

FIRST AUTHOR	AGE OF PATIENT/SEX	MUSCLE SYMPTOMS	CREATINE KINASE	ILD	SKIN ULCERS	TREATMENT BEFORE SPnM	TREATMENT AFTER PnM	TIME TO RESOLUTION OF SPnM	OUTCOME
Powell ⁶	34/M	No	Normal	Yes	No	PNS + AZA	Pulse IV methyl-PNSL 2g, PNS 80mg/day	N/A	Deceased
Bradley ¹³	42/M	Yes	Elevated	Yes	Yes	PNS 80mg/day + AZA 100mg/day	Same as before	1 month	Alive
Yoshida ¹⁷	38/M	Yes	Elevated	No	No	PNSL 40mg/day	3 weekly pulses of CPA 50mg + CSPN 300mg + PNSL 50mg/day + MTX 7.5mg/week	3 months	Alive
de Souza Neves ¹⁸	45/M	Yes	Normal	Yes	Yes	none	PNSL 1mg/kg + chloroquine diphosphate 250mg/day + MTX 20mg + IV gamma globulins + pulse CPA 1g	2 months	Alive
Korkmaz ²⁰	28/M	Yes	Normal	Yes	No	PNSL 60mg/day	IV CPA 1g/month + PNSL 50mg/day tapered to 10mg/day after 4 weeks	3 months	Alive
Kim ²¹	38/F	Yes	Elevated	Yes	Yes	PNSL 30mg/day + hydroxy-chloroquine 300mg/day	PNSL 30mg/day + hydroxychloroquine 300mg/day + CSPNA 100mg/day	1 month	Alive
Kuroda ²²	46/M	Yes	Elevated	Yes	No	PNSL 60mg/day	PNSL taper + CSPNA 100mg/day	4 months	Alive
Barvaux ²³	42/M	Yes	N/A	Yes	No	PNSL 0.5mg/kg + MTX 15mg/week + hydroxy-chloroquine 400mg/day	5 monthly IV pulses of CPA (0.75 to 1.5g/m ²) + IV gamma globulins (0.4g/kg for 5 consecutive days + single dose every 6 weeks	12 months +	Alive
Marsrouha ²⁵	66/M	N/A	Normal	Yes	No	PNS + MTX	Pulse steroid + MTX	4 months	Alive
Lee ²⁶	25/F	Yes	Elevated	No	No	PNSL 50mg/day + MTX 10mg/week	1g IV rituximab on Days 1 and 14, 740mg IV CPA + 500mg IV methyl-PNSL on Days 2 and 15	Several weeks	Alive
Terao ²⁷	16/M	Yes	Normal	Yes	Yes	Topical steroids	Bed rest, followed by PNSL 40mg/day + CSPNA 125mg/day	2 months	Alive
Current case	52/F	No	Normal	Yes	Yes	PNS 60mg/day + hydroxy-chloroquine 200mg twice/day	CSPNA 100mg twice/day, hydroxychloroquine 200mg/twice per day, PNS 30mg/day	1 week	Alive

ILD=interstitial lung disease; PnM=pneumomediastinum; PNS=prednisone; PNSL=prednisolone; AZA=azathioprine; CSPNA=cyclosporine A; CPA=cyclophosphamide; MTX=methotrexate, N/A=not available

REFERENCES

1. Dalakas MC, Hohlfeld R. Polymyositis and dermatomyositis. *Lancet*. 2003;362:971–982.
2. Mascaro JM, Hausmann G, Herrero C, et al. Membrane attack complex deposits in cutaneous lesions of dermatomyositis. *Arch Dermatol*. 1995;131:1386–1392.
3. Hill C, Zhang Y, Sigurgeirsson B, et al. Frequency of specific cancer types in dermatomyositis and polymyositis: a population-based study. *Lancet*. 2001;357:96–100.
4. András C, Ponyi A, Constantin T, et al. Dermatomyositis and polymyositis associated with malignancy: a 21-year retrospective study. *J Rheumatol*. 2008;35:438–444.
5. Dickey B, Myers A. Pulmonary disease in polymyositis and dermatomyositis. *Semin Arthritis Rheumatol*. 1984;41:137–140.
6. Powell C, Kendall B, Wernick R, Heffner J. A 34-year-old man with amyopathic dermatomyositis and rapidly progressive dyspnea with facial swelling. *Chest*. 2007;132:1710–1713.
7. Hochberg M, Feldman D, Stevens M, et al. Antibody to Jo-1 in polymyositis/dermatomyositis association with interstitial pulmonary disease. *J Rheumatol*. 1984;11:663–665.
8. Bernstein R, Morgan S, Chapman J, et al. Anti Jo-1 antibody a marker for myositis with interstitial lung disease. *Br Med J*. 1984;289:152–155.
9. Phillips T, Leigh I, Wright J. Dermatomyositis and pulmonary fibrosis associated with anti-Jo-1 antibody. *J Am Acad Dermatol*. 1987;17:381–382.
10. High W, Cohen J, Murphy B, Costner M. Fatal interstitial pulmonary fibrosis in anti-Jo-1-negative amyopathic dermatomyositis. *J Am Acad Dermatol*. 2003;49:295–298.
11. Morganroth P, Kreider M, Okawa J, et al. Interstitial lung disease in classic and skin-predominant dermatomyositis. *Arch Derm*. 2010;146:729–737.
12. Mukae H, Ishimoto H, Sakamoto N, et al. Clinical differences between interstitial lung disease associated with clinically amyopathic dermatomyositis and classic dermatomyositis. *Chest*. 2009;136:1341–1347.
13. Bradley J. Spontaneous pneumomediastinum in adult dermatomyositis. *Ann Rheum Dis*. 1986;45:780–782.
14. Jansen T, Barrera P, van Engelen B, et al. Dermatomyositis with subclinical myositis and spontaneous pneumomediastinum with pneumothorax: case report and review of the literature. *Clin Exp Rheumatology*. 1998;16:733–735.
15. Cicuttini F, Fraser K. Recurrent pneumomediastinum in adult dermatomyositis. *J Rheum*. 1987;16:384–386.
16. Kono H, Inokuma S, Nakayama H, Suzuki M. Pneumomediastinum in dermatomyositis: association with cutaneous vasculopathy. *Ann Rheum Disease*. 2000;59:361–372.
17. Yoshida K, Kurosaka D, Kingetsu I, et al. Pneumomediastinum in dermatomyositis itself is not a poor prognostic factor: report of a case and review of the literature. *Rheumatol Int*. 2008;28:913–917.
18. Neves F, Shinjo S, Carvalho J, et al. Spontaneous pneumomediastinum and dermatomyositis may be a not so rare association: report of a case and review of the literature. *Clin Rheum*. 2007;26:105–107.
19. Yamanishi Y, Maeda H, Konishi F, et al. Dermatomyositis associated with rapidly progressive fatal interstitial pneumonitis and pneumomediastinum. *Scan J Rheumatol*. 1999;28:58–61.
20. Korkmaz C, Ozkan R, Akay M, Hakan T. Pneumomediastinum and subcutaneous emphysema associated with dermatomyositis. *Rheumatology*. 2001;40:476–478.
21. Kim HJ, Hong YK, Yoo WH. Dermatomyositis, complicated with pneumomediastinum, successfully treated with cyclosporine A: a case report and review of the literature. *Rheumatol Int*. 2009;29:1101–1104.
22. Kuroda T, Morikawa H, Satou T, et al. A case of dermatomyositis complicated with pneumomediastinum successfully treated with cyclosporine A. *Clin Rheumatol*. 2003;22:45–48.
23. Barvaux V, Mullem X, Pieters T, Houssiau F. Persistent pneumomediastinum and dermatomyositis: a case report and review of the literature. *Clin Rheumatol*. 2001;20:359–361.
24. Ozawa Y, Kurosaka D, Yokoyama T, Tajima N. Therapeutic efficacy of cyclosporin A in four cases of amyopathic dermatomyositis with rapidly progressive interstitial pneumonia. *Ryumachi*. 2000;40:798–809.
25. Masrouha K, Kanj N, Uthman I. Late-onset pneumomediastinum in dermatomyositis. *Rheumatol Int*. 2009;30:291–292.
26. Lee M, Hutchinson D. Spontaneous pneumomediastinum secondary to refractory dermatomyositis successfully treated with rituximab. *Clin Rheumatol*. 2010. [Epub ahead of print].
27. Terao M, Ozawa K, Inui S, et al. A case of dermatomyositis complicated with pneumomediastinum. *Mod Rheumatol*. 2007;17:156–159. ●